

ORIGINAL ARTICLE

Traditional grazing management creates heterogeneous swards and fosters grasshopper densities

Cinja Schwarz  and Thomas Fartmann 

Department of Biodiversity and Landscape Ecology, Faculty of Biology and Chemistry, Osnabrück University, Osnabrück, Germany

Abstract Common pastures were once the dominant type of land use in many European regions. However, during the past 150 years, they have declined dramatically. Recent studies have shown that they are hotspots for rare plant, butterfly, and bird species in the study area, the Bavarian pre-Alps (southern Germany). However, studies on the value of these pastures for Orthoptera (hereinafter termed “grasshoppers”) have been scarce. Here, we studied the effects of traditional summer grazing in common pastures on grasshopper assemblages. Our study revealed that grasshopper species richness did not differ between common pastures ($n = 57$) and controls ($n = 57$). By contrast, density of all and of threatened species varied between common pastures and controls in all plots and within the two vegetation types with the highest grasshopper abundance, grasslands on mineral soil and fens. Two threatened species, *Pseudochorthippus montanus* and *Stethophyma grossum*, were identified as indicators for common pastures; controls had no indicative species. Traditional low-intensity grazing in common pastures has resulted in open and heterogeneous swards with some bare ground, a low cover of litter and an intermediate vegetation height favouring high densities of grasshopper species in general and threatened species in particular. This is especially true for the two most productive vegetation types, grasslands on mineral soil and fens. To promote biodiversity in general and grasshopper densities in particular, we recommend maintaining traditional cattle grazing (stocking capacities: 0.5–2.0 livestock units/ha) in common pastures. Where possible, this grazing regime should also be introduced in the surrounding landscape.

Key words biodiversity conservation; common pasture; habitat heterogeneity; insect abundance; land-use change; mire ecosystem

Introduction

Since the beginning of the industrial era 200 years ago, humans have altered their environment at an unprecedented rate (Foley *et al.*, 2005; Rockström *et al.*, 2009). As a result, biodiversity is in a steep decline and scientists suspect that we are heading for a 6th mass extinction (Barnosky *et al.*, 2011; Dirzo *et al.*, 2014). The loss of biodiversity jeopardizes ecosystem functioning and our

own well-being (Dirzo *et al.*, 2014; Ripple *et al.*, 2017; IPBES, 2019; Cardoso *et al.*, 2020). Hence, halting the decline is one of the major challenges for humanity in the present time.

Insects are the most species-rich taxonomic group worldwide (Stork, 2018). However, their decline is much stronger than those of plants or vertebrates (Thomas *et al.*, 2004; Sánchez-Bayo & Wyckhuys, 2019; Cardoso *et al.*, 2020; Wagner, 2020). The loss of insects has cascading effects on various other organisms within ecosystems (Cardoso *et al.*, 2020; Wagner, 2020). For example, insect prey abundance directly affects population size of insectivorous species of higher trophical levels (*e.g.*, birds) (Fartmann *et al.*, 2021). As for biodiversity in general, land-use change has been identified as the main

Correspondence: Cinja Schwarz, Department of Biodiversity and Landscape Ecology, Faculty of Biology and Chemistry, Osnabrück University, Barbarastraße 11, 49076 Osnabrück, Germany. Email: cinja.schwarz@uos.de

driver of the decline of insects (IPBES, 2019; Cardoso *et al.*, 2020; Wagner, 2020).

Nutrient-poor, semi-natural grasslands are among the most species-rich ecosystems across Europe (Dengler *et al.*, 2014; Chytrý *et al.*, 2015; Feurdean *et al.*, 2018). However, due to the transition from pre-industrial land use to modern-day agricultural practices, the extent of such grasslands has greatly decreased and the remaining patches have often suffered from habitat deterioration, mainly due to land-use intensification or succession as a consequence of land-use abandonment (Wallis de Vries & Poschlod, 2002; Poschlod & Braun-Reichert, 2017; Fartmann *et al.*, 2021). Since 1992, some types of traditionally managed semi-natural grasslands and some of their characteristic species have become protected by the Habitats Directive (EC, 2007). However, still several types of grasslands with high value for insect conservation, for example, wet grasslands on mineral soil or different types of nutrient-poor pastures, have not been covered appropriately by the directive (Gepp, 2001; Plieninger *et al.*, 2015). Additionally, among the over 1 000 species protected by the directive, only 70 are insects (Dirzo *et al.*, 2014). This bias can at least partly be explained by the limited number of studies on insect declines and the minimal attention paid to the loss of insect diversity at the time.

Until the mid-19th century, common pastures were the dominant type of land use in many European regions and had a high economic value (Lederbogen *et al.*, 2004; Brown, 2006; Plieninger *et al.*, 2006; Hrvatin & Perko, 2008). However, since then, they have declined dramatically in area and economic importance. The pre-Alps in southern Bavaria (Germany) are one of the last strongholds for common pastures in Central Europe (Lederbogen *et al.*, 2004; Thumm *et al.*, 2005). Here, the common pastures comprise mosaics of mire ecosystems, grasslands on mineral soil, and light woodlands (Schwarz *et al.*, 2018). Recent studies have shown that these pastures are hotspots for rare plant, butterfly, and bird species (Anthes *et al.*, 2003; Rosenthal & Lederbogen, 2008; Weking *et al.*, 2013; Schwarz *et al.*, 2018; Schwarz & Fartmann, 2021). However, studies on the value of these pastures for Orthoptera (hereinafter termed “grasshoppers”) have been scarce. Grasshoppers have a high functional significance in grasslands due to their key role as herbivores and prey (Samways, 2005). Moreover, they rapidly respond to changes in land use (Marini *et al.*, 2008; Bazelet & Samways, 2012; Uchida & Ushimaru, 2014; Fumy *et al.*, 2021). Hence, they are well-established bioindicators (Samways, 2005; Sliacka *et al.*, 2013; Sergeev, 2021).

Here, we studied the effects of traditional summer cattle grazing in common pastures of the Bavarian pre-Alps on grasshopper assemblages. Therefore, we compared environmental conditions and assemblage composition (species richness, density, indicator species) between common pastures ($n = 57$) and surrounding grasslands on mineral soil and open mires as a control ($n = 57$). In particular, we addressed the following research questions:

(i) How do environmental conditions as well as grasshopper species richness, density, and indicator species vary between common pastures and controls?

(ii) What are the drivers of grasshopper assemblage composition?

(iii) Which implications for grasshopper conservation can be drawn from our findings?

Materials and methods

Study area

The study area (hereafter referred to as pre-Alps) is located at the northern foothills of the Alps (750–900 m a.s.l.) in the south of the German Federal State of Bavaria. The landscape was formed by the last glaciation and is characterized by a heterogeneous young moraine relief (Bayerisches Geologisches Landesamt, 1996). Due to a rather cool and wet climate (mean annual temperature: 7.7 °C; mean annual precipitation: 1 336 mm; meteorological station Bad Kohlgrub [742 m a.s.l.]; period: 1992–2019; DWD, 2020), numerous mires developed in the valleys and hollows (Succow & Jeschke, 1990; Bayerisches Geologisches Landesamt, 1996). Today, the study area contains one of the largest networks of mire ecosystems in Central Europe with an outstanding value for biodiversity conservation and, hence, belongs to one of the German biodiversity hotspots (Ackermann *et al.*, 2012; BfN, 2012).

Grasslands, mainly used for dairy farming, are the dominant type of land use in the pre-Alps (BfN, 2012). However, besides large areas of improved grasslands, traditionally managed grasslands such as common pastures, litter meadows (meadows mown in autumn to obtain bedding for livestock), and hay meadows have remained due to the geographical, climatical, and economical marginalization of the study area (Lederbogen *et al.*, 2004; Brown, 2006; Schwarz & Fartmann, 2021).

Sampling design

Plots We studied seven randomly selected common pastures, which are traditionally managed by rural




Fig. 1 Location of the study area and common pastures in Upper Bavaria (southern Germany). Common pasture: BS = Berghofer Söldner; EG = Echelsbach Gschwend; HA = Hachegger Viehweide; MO = Moosreitener Viehweide; MV = Mühlenberger Viehweide; PV = Premer Viehweide; RV = Rieder Viehweide.

communities, and their adjacent grasslands and mires (Fig. 1). Each common pasture with its adjacent grasslands and mires is hereafter referred to as a “subarea.” Common pastures had a size of 5–121 ha (mean \pm SE: 57 ± 15). Grazing was performed with brown dairy cattle in low stocking capacities (0.5–2.0 livestock units [LSU]/ha) from May to October (Lederbogen *et al.*, 2004; Thumm *et al.*, 2005, own observation). Overall, common pastures were characterized by a high heterogeneity of habitat types. In addition to large areas of grasslands on mineral soil, the pastures comprised mosaics of grazed fens, transition mires, and raised bogs (Schwarz *et al.*, 2018). In all seven common pastures, low-intensity grazing has been conducted for centuries. Except grasslands on mineral soils, the common pastures have largely never been fertilized (Lederbogen *et al.*, 2004).

To compare common pastures with ungrazed surrounding grasslands on mineral soil and open mires, we used a paired design. Therefore, we randomly selected 57 plots with a size of 500 m² (20 m \times 25 m) in common pas-

tures. To contrast data from the common pastures with those of the adjacent grasslands on mineral soil and open mires, 57 control plots with the same size were randomly chosen. Each plot had a distance between 20 and 25 m to the fence of the respective common pasture to avoid edge effects (Schirmel *et al.*, 2010). Altogether, we divided the plots into the following four vegetation types using character plant species according to Oberdorfer (1992) and Dierßen & Dierßen (2008): (i) grasslands on mineral soil, (ii) fens, (iii) transition mires, and (iv) raised bogs (Table 1) (Schwarz & Fartmann, 2021). Additionally, land use of control plots was differentiated into the three land-use types: (i) fallow, (ii) mown once, and (iii) mown twice or more (Table 1).

Habitat quality To assess habitat quality, we sampled several environmental variables once per plot in August 2015 (Table 2). Cover of different vegetation layers was estimated for a randomly selected subplot of 9 m² (3 m \times 3 m), except the cover of shrubs, which was estimated

Table 1 Absolute and relative frequencies of the four vegetation types in common pastures ($n = 57$) and controls ($n = 57$) as well as of the three land-use types in controls (abandoned: $n = 28$; mown once: $n = 24$; mown twice or more often: $n = 5$). Differences in absolute frequencies were tested using Chi-squared test. Common pastures vs. control: not significant; land-use types of the control plots: $P \leq 0.001$.

Vegetation type	Common pasture		Control		Control					
	<i>n</i>	%	<i>n</i>	%	Abandoned		Mown once		Mown \geq twice	
					<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Grassland on mineral soil	14	25	13	23	1	4	7	29	5	100
Fen	9	16	12	21	4	14	8	33	0	0
Transition mire	17	30	15	26	6	21	9	38	0	0
Raised bog	17	30	17	30	17	61	0	0	0	0

for the whole plot. Vegetation density was determined in 0–5 cm, 10–15 cm and 25–30 cm height from ground using a wire-frame box of 30 cm depth and 50 cm width (Poniatowski & Fartmann, 2008; Löffler & Fartmann, 2017). Vegetation height was measured at three random points in each plot and averaged for further analysis. We measured the potential sunshine duration (accuracy 0.5 h) using a horizontoscope after Tonne (Scherer *et al.*, 2021) for the month of August in the center of each plot. Cow droppings were counted in every pasture plot on a randomly selected subplot of 100 m² (10 m × 10 m) as a proxy for grazing intensity.

Grasshopper assemblages We sampled grasshopper assemblages in August 2015, when nearly all species are adult (Schlumprecht & Waeber, 2003). Grasshopper densities were investigated once for each plot using a box quadrat (1.41 × 1.41 m) (Gardiner *et al.*, 2005). The quadrat was placed randomly 10 times per plot (total area = 20 m²), and individuals of each species within the quadrat were counted (Fartmann *et al.*, 2008; Poniatowski & Fartmann, 2010; Helbing *et al.*, 2014). Grasshoppers were determined using Bellmann (2006) and nomenclature follows Fischer *et al.* (2020). For further analysis, we classified species as either threatened (including near-threatened species) or non-threatened according to the current red data book of Bavaria (Voith *et al.*, 2016).

Statistical analysis

We performed a Chi-squared test to examine differences in the nominal variables vegetation type between common pastures and land-use type within controls

(Table 1). Differences in all metric environmental variables, species richness, and density of grasshoppers (response variables) between common pastures and controls (predictor variable) were tested using a generalized linear mixed-effects model (GLMM) with “subarea” as a random factor and an observation-level random intercept nested in “subarea” in case of overdispersion (error structure: binomial, negative binomial, Poisson, or Gaussian, respectively) (Table 2 and Fig. 2). All pairwise comparisons between land-use types were made using the function “glht()” in the multcomp package (Hothorn *et al.*, 2008). Moreover, we conducted an indicator species analysis (ISA) (Dufrene & Legendre, 1997) to identify species indicative for common pastures and controls in general and within each of the four vegetation types (Table 3).

To ascertain those environmental variables (predictor variables) that explain grasshopper species richness and density (response variables) in common pastures best, further GLMMs with “subarea” as a random factor were conducted. Model averaging based on an information-theoretic approach was used to increase model robustness and identify the most important variables (Burnham & Anderson, 2002; Grueber *et al.*, 2011). We applied the “dredge” function (R package MuMIn, Bartón, 2016) and only included top-ranked models within $\Delta AIC_C < 3$ for model averaging (*cf.* Grueber *et al.*, 2011). Ranges of marginal and conditional R^2 values were calculated as measures of the explanatory power of the top-ranked models (Nakagawa & Schielzeth, 2013; Nakagawa *et al.*, 2017). We calculate the variance inflation factor (VIF) and used only variables with $VIF \leq 5$ in the GLMMs to avoid multicollinearity (Sheather, 2009). Statistical analysis was performed using R 3.4.3.

Table 2 Mean values (\pm standard errors) of environmental variables in common pastures ($n = 57$) and controls ($n = 57$) in general and within the four vegetation types (grassland on mineral soil: $n = 27$; fen: $n = 21$; transition mire: $n = 32$; raised bog: $n = 34$). Differences were tested using Generalized Linear Mixed-Effects Models (GLMM). In all models, ‘subarea’ was set up as random factor. In the case of overdispersion, an observation level random effect was used. Bold values: Variable differs significantly. Error structures were marked with superscripted letters after P values: b = binomial, nb = negative binomial, p = Poisson, g = Gaussian. n.s. = not significant, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

Variable	Grassland on mineral soil			Fen			Transition mire			Raised bog					
	Pasture	Control	P	Pasture	Control	P	Pasture	Control	P	Pasture	Control	P			
Cover (%) of															
Shrubs	6.6 \pm 1.4	7.4 \pm 1.6	n.s. ^b	0.4 \pm 0.4	0.0 \pm 0.0	n.s. ^b	1.1 \pm 0.7	4.2 \pm 2.5	n.s. ^b	7.9 \pm 3.2	2.7 \pm 1.9	** ^b	13.2 \pm 2.5	19.6 \pm 3.4	n.s. ^b
Poaceae	19.8 \pm 2.5	23.1 \pm 3.0	n.s. ^b	37.9 \pm 4.2	41.5 \pm 6.6	n.s. ^b	18.3 \pm 5.5	24.4 \pm 6.3	n.s. ^b	21.6 \pm 4.6	28.0 \pm 4.7	n.s. ^b	4.0 \pm 0.6	3.7 \pm 0.4	n.s. ^b
Cyper-aceae	52.8 \pm 2.3	48.8 \pm 2.8	n.s. ^b	42.9 \pm 3.8	30.8 \pm 6.7	n.s. ^b	68.9 \pm 6.0	54.6 \pm 6.9	* ^b	52.9 \pm 4.8	55.0 \pm 3.9	n.s. ^b	52.4 \pm 2.9	52.9 \pm 3.4	n.s. ^b
Litter	16.1 \pm 1.7	15.0 \pm 1.9	n.s. ^b	10.4 \pm 2.5	8.8 \pm 1.3	n.s. ^b	14.4 \pm 3.9	10.4 \pm 2.2	n.s. ^b	16.8 \pm 3.1	16.8 \pm 5.9	n.s. ^b	21.2 \pm 3.4	21.2 \pm 2.9	n.s. ^b
Mosses	61.6 \pm 3.8	55.3 \pm 4.3	n.s. ^b	36.3 \pm 7.5	31.7 \pm 8.5	n.s. ^b	68.3 \pm 7.5	51.7 \pm 7.2	n.s. ^b	58.2 \pm 4.9	45.0 \pm 8.2	n.s. ^b	82.4 \pm 5.5	84.9 \pm 3.6	n.s. ^b
Open soil	5.3 \pm 0.7	2.4 \pm 0.5	*** ^b	5.5 \pm 1.3	2.1 \pm 0.8	* ^b	3.1 \pm 1.1	2.7 \pm 1.4	n.s. ^b	8.4 \pm 1.9	4.7 \pm 0.9	*** ^b	3.1 \pm 0.7	0.4 \pm 0.2	*** ^b
Vegetation height (cm)	21.8 \pm 1.4	23.1 \pm 2.1	n.s. ^{nb}	23.5 \pm 3.9	22.0 \pm 3.8	n.s. ^{nb}	24.8 \pm 2.7	22.5 \pm 3.6	** ^g	24.1 \pm 2.5	23.6 \pm 2.4	** ^{nb}	16.5 \pm 0.9	24.0 \pm 5.8	n.s. ^{nb}
Vegetation density (%)															
0–5 cm height	91.4 \pm 1.7	93.7 \pm 1.2	n.s. ^b	96.3 \pm 2.2	98.7 \pm 0.8	n.s. ^b	97.8 \pm 0.9	94.2 \pm 2.0	n.s. ^b	95.9 \pm 1.5	96.0 \pm 1.6	n.s. ^b	79.4 \pm 3.7	87.6 \pm 3.0	** ^b
25–30 cm height	6.5 \pm 1.1	11.4 \pm 2.7	n.s. ^b	5.9 \pm 1.3	15.2 \pm 7.8	*** ^b	4.7 \pm 1.7	8.5 \pm 3.8	n.s. ^b	9.4 \pm 3.1	16.8 \pm 6.6	n.s. ^b	5.0 \pm 1.5	5.6 \pm 2.1	n.s. ^b
Sunshine duration (h)	11.5 \pm 0.3	11.6 \pm 0.2	n.s. ^p	11.4 \pm 0.6	11.0 \pm 0.6	n.s. ^{nb}	10.8 \pm 0.8	12.1 \pm 0.3	n.s. ^{nb}	11.0 \pm 0.7	11.9 \pm 0.3	n.s. ^{nb}	12.3 \pm 0.2	11.5 \pm 0.4	n.s. ⁿ




Fig. 2 Mean values (\pm standard error) of the number of all species (A), number of threatened species (B), density of all species (C), and density of threatened species (D) in common pastures ($n = 57$) and controls ($n = 57$). Differences were tested using GLMM. In all models, “subarea” was set up as random factor. In the case of overdispersion, an observation level random effect was used. n.s. = not significant, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

Results

Plots

The composition of vegetation types did not differ between common pastures and controls (Table 1). Grasslands on mineral soil, fens, transition mires and raised bogs always accounted for a share of around one quarter of the plots. By contrast, land use clearly varied between common pastures and controls. All common pastures were grazed ($n = 57$) (cf. Section Sampling design) whereas one half of the controls was abandoned, two fifths were mown once (usually used as litter meadows) and the remaining tenth comprised grasslands mown twice or more (Table 1). Within the controls, land-use intensity decreased from grasslands on mineral soil to fens/transition mires to raised bogs.

Land use had a clear effect on environmental conditions on the plots (Table 2). Grazing intensity in common pastures, indicated by the density of cow droppings, decreased in the following order: grasslands on mineral soil (mean no. of cow droppings/100 $m^2 \pm$ SE: 14.1 ± 2.3), fens (6.8 ± 1.8), transition mires (3.2 ± 0.8), and raised bogs (1.6 ± 0.6). Cow-dropping density differed statistically between grasslands on mineral soil and those of the three other vegetation types (GLMM, $P \leq 0.05$). Cattle grazing in common pastures compared to controls resulted in a higher cover of open soil (all plots as well as plots of grasslands on mineral soil, transition mires, and raised bogs) and a lower vegetation density (plots of grasslands on mineral soil and raised bogs). Additionally, within fens, common pastures had a higher cover of Cyperaceae and a taller vegetation than controls. Moreover, within transition mires used as common

Table 3 Results of ISA (Dufrêne & Legendre, 1997) for common pastures and controls in general and within the four vegetation types. Threat status (TS): threatened species in Bavaria (Voith *et al.*, 2016). Values hatched in grey: species that are indicator species for this land-use type. ab = relative abundance, fr = frequency, n.s. = not significant, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

Species	TS	P	Mineral soil						Fen						Transition mire						Raised bog					
			Pasture			Control			Pasture			Control			Pasture			Control			Pasture			Control		
			ab	fr	ab	fr	ab	fr	ab	fr	ab	fr	ab	fr	ab	fr	ab	fr	ab	fr	ab	fr	ab	fr		
<i>Ch. albo-marginatus</i>	.	n.s.	43	11	57	9	35	29	51	31	0	0	9	8	5	12	0	0	0	0	0	0	0			
<i>Ch. brunneus</i>	.	n.s.	0	0	100	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	100	6			
<i>Ch. dorsatus</i>	x	n.s.	78	47	22	46	53	100	18	92	24	78	2	67	1	29	1	33	0	6	0	6	6			
<i>Chr. dispar</i>	.	n.s.	0	0	100	2	0	0	0	0	0	0	0	0	0	0	100	7	0	0	0	0	0			
<i>C. fuscus</i>	.	n.s.	49	42	51	46	23	64	16	54	18	67	25	92	8	53	10	53	0	0	0	0	0			
<i>D. verrucivorus</i>	x	n.s.	0	0	100	4	0	0	0	0	0	0	100	17	0	0	0	0	0	0	0	0	0			
<i>E. brachyptera</i>	.	n.s.	33	12	67	23	25	29	17	23	10	22	11	42	1	6	37	33	0	0	0	0	0			
<i>M. brachyptera</i>	x	n.s.	40	53	60	39	1	7	7	8	12	56	5	17	9	53	8	27	20	88	39	88				
<i>O. viridulus</i>	x	n.s.	36	5	64	9	14	7	45	23	0	0	24	17	17	12	0	0	0	0	0	0	0			
<i>P. montanus</i>	x	**	61	98	39	89	17	100	9	77	30	100	14	100	11	94	11	100	5	100	3	82				
<i>P. parallelus</i>	.	n.s.	65	9	35	5	60	29	37	23	0	0	0	0	4	6	0	0	0	0	0	0	0			
<i>R. roeselii</i>	.	n.s.	53	14	47	14	46	43	19	23	5	11	21	25	1	6	8	13	0	0	0	0	0			
<i>S. grossum</i>	x	*	67	40	33	16	15	21	22	23	2	22	8	8	25	53	7	20	20	53	1	12				
<i>T. subulata</i>	.	n.s.	61	18	39	12	49	57	16	23	11	22	21	25	0	0	0	0	0	0	0	3	6			
<i>T. undulata</i>	.	n.s.	0	0	100	7	0	0	0	0	0	0	53	17	0	0	0	0	0	0	0	47	12			
<i>T. cantans</i>	.	n.s.	36	5	64	9	9	7	0	0	0	0	43	25	15	6	17	7	8	6	8	6	6			




Fig. 3 Relationship between the number of all species (mean \pm SE: 3.5 ± 0.2), density of all species (15.9 ± 2.3), and density of threatened species (14.0 ± 2.0) and environmental variables in common pastures (see Table 4 for detailed GLMM statistics).

pastures, the cover of shrubs and the vegetation height were higher.

Grasshopper assemblages and response to environmental conditions

Overall, we recorded 2 891 grasshopper individuals belonging to 16 species (Table 3). Five of these species

are considered threatened. Species richness of all and of threatened species did not differ between common pastures and controls (all plots and within all four vegetation types; Fig. 2A and B). By contrast, density of all and of threatened species varied in all plots and within the 2 vegetation types with the highest grasshopper abundance, grasslands on mineral soil and fens, between common pastures and controls (Fig. 2C and D).

Table 4 Model-averaging results (GLMM) of the relationship between the response variables number of all species (Poisson) (A), number of threatened species (Poisson) (B), density of all species (negative binomial) (C), and density of threatened species (negative binomial) (D) and environmental variables (predictor variables) in common pastures. Differences were tested using Generalized Linear Mixed-Effects Models (GLMM). In all models, ‘subarea’ was set up as random factor. Predictor variables tested in all models were cover of shrubs, Cyperaceae, Poaceae, herbs, litter, mosses, open soil, vegetation height, vegetation density (25–30 cm height) and sunshine duration. For all models, model-averaged coefficients (full average) were derived from the top-ranked model (Δ AICc < 3). The explanatory power of the models is shown by ranges of marginal and conditional R^2 values (Nakagawa & Schielzeth, 2013; Nakagawa *et al.*, 2017) of the top-ranked models (Δ AICc < 3). n.s. = not significant, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

All species					Threatened species				
Variable	Est.	SE	Z	P	Variable	Est.	SE	Z	P
(A) No. species					(B) No. species				
(Intercept)	1.39	0.08	16.92	***	(Intercept)	–	–	–	–
Shrubs	–0.02	0.01	–2.51	*	Shrubs	–	–	–	–
$R^2_M = 0.16, R^2_C = 0.16$									
(C) Density					(D) Density				
(Intercept)	2.14	0.15	14.54	***	(Intercept)	2.13	0.49	4.30	***
Cyperaceae	0.03	0.01	6.31	***	Cyperaceae	0.03	0.01	3.90	***
Litter	–0.04	0.01	3.56	***	Litter	–0.04	0.01	2.75	**
Poaceae	0.03	0.01	5.74	***	Poaceae	0.03	0.01	4.08	***
Shrubs	–0.01	0.01	0.83	n.s.	Shrubs	–0.01	0.01	0.67	n.s.
$R^2_M = 0.33–0.38, R^2_C = 0.57–0.67$					$R^2_M = 0.36–0.40, R^2_C = 0.51–0.61$				

Considering all plots, two threatened species, *Pseudochorthippus montanus* and *Stethophyma grossum*, were identified as indicators of common pastures (Table 3). By contrast, controls had no indicative species. Within three of the four vegetation types, further indicator species were detected. In grasslands on mineral soil, *Chorthippus dorsatus*, *Pseudochorthippus parallelus*, *Roeseliana roeselii*, and *Tetrix subulata* were ascertained as indicator species of common pastures. The former is a threatened species. By contrast, within this vegetation type, *Chorthippus albomarginatus* was indicative for controls. In fens, the threatened species *P. montanus* was an indicator for common pastures and *Conocephalus fuscus* and the threatened species *Decticus verrucivorus* were characteristic for controls. Additionally, in raised bogs, *Metrioptera brachyptera* was identified as an indicator species of controls.

The GLMM analysis revealed that a higher cover of shrubs had negative effects on overall species richness in common pastures (Fig. 3A, Table 4). The cover of shrubs was negatively correlated with the number of cow droppings and vegetation density in 0–5 cm height (Table S1). The GLMM analysis for the number of threatened species failed to identify environmental predictors (Table 4). By contrast, GLMM analyses on the drivers of density of all and of threatened species in common pastures produced very similar results and had a high explanatory power

(Fig. 3B–G, Table 4). Grasshopper density increased with the cover of grasses (Poaceae and Cyperaceae) but decreased with the cover of litter and shrubs. The cover of Poaceae was positively correlated with the number of cow droppings and vegetation density in 0–5 cm height (Table S1). Values of marginal R^2 ranged between 0.33 and 0.40.

Discussion

Our study revealed that grasshopper species richness did not differ between common pastures and controls in the Bavarian pre-Alps. However, density of all and of threatened species varied between common pastures and controls in all plots and within the two vegetation types with the highest grasshopper abundance, grasslands on mineral soil and fens. Two threatened species, *P. montanus* and *S. grossum*, were identified as indicators for common pastures. By contrast, controls had no indicative species.

Usually, changes in abundance of species reflect alterations in environmental conditions much earlier than shifts in species richness (Wettstein & Schmid, 1999; Fartmann *et al.*, 2012; Löffler & Fartmann, 2017) since declines in the latter can only be detected if a population has completely disappeared (Thomas *et al.*, 2006; Dirzo *et al.*, 2014). Composition of vegetation types

was very similar between common pastures and controls; solely land use varied. Hence, we observed differences in grasshopper density but not in species richness between common pastures and controls.

The key variables determining habitat quality for grasshoppers are (i) a suitable (warm) microclimate, (ii) enough food resources, (iii) appropriate oviposition sites, and (iv) refuges against predators or adverse weather (Gardiner & Dover, 2008; Wunsch *et al.*, 2012). Such diverse requirements are usually only fulfilled in habitats that are shaped by some kind of low-intensity land use (Kruess & Tschardt, 2002; Löffler & Fartmann, 2017; Fumy *et al.*, 2021). Low-intensity grazing in common pastures has led to open and heterogeneous vegetation with some bare ground, a low cover of litter and an intermediate vegetation height favouring high densities of grasshoppers in general and threatened species in particular. The two threatened indicator species of common pastures, *P. montanus* and *S. grossum*, are also known to achieve their highest abundance in open wet grasslands and mires with low management intensity (Schlumprecht & Waeber, 2003).

Grasshopper species richness within common pastures was favored by a low cover of shrubs. Shrub cover reflected grazing intensity within the pastures since it was negatively correlated with the density of cow droppings. Grazing intensity in turn was related to the productivity of the vegetation types (*cf.* Lederbogen *et al.*, 2004; Ellenberg & Leuschner, 2010) and, hence, decreased in the order: grasslands on mineral soil, fens, transition mires, and raised bogs. Undergrazing favors the establishment of shrubs and taller vegetation rich in litter (Löffler *et al.*, 2019; Fumy *et al.*, 2021). As a result, patches of sunlit bare ground, vital for soil-breeding species, disappear and microclimatic conditions become cooler (Stoutjesdijk & Barkman, 1992), with negative effects on grasshopper species (Marini *et al.*, 2008; Helbing *et al.*, 2014). By contrast, for the number of threatened species within common pastures, the GLMM failed to identify environmental predictors, which we attribute to the relatively constant number of threatened species across the environmental gradient.

However, the GLMM analyses on the drivers of density of all and of threatened species in common pastures produced consistent results and had a high explanatory power. Density of both groups increased with the cover of grasses (Poaceae and Cyperaceae) but decreased with the cover of litter and shrubs. The majority of grasshopper species feed on grasses (Ingrisch & Köhler, 1998). Recent studies have shown that the food supply may limit grasshopper densities in habitats with a sparse vegetation (Helbing *et al.*, 2014; Löffler *et al.*, 2016). In the com-

mon pastures, this seems to be especially true for raised bogs where the cover of grasses was much lower than in grasslands on mineral soil or in fens. Additionally, undergrazing, indicated by a higher cover of litter and shrubs, had not only negative effects on species richness but also on density of grasshoppers.

In contrast to common pastures, only two fifths of the control plots were characterized by low-intensity management, that is, meadows mown once, usually used as litter meadows. One half of the controls were abandoned and the remaining tenth comprised grasslands mown twice or more. Both abandonment and intensive land use have negative impacts on grasshoppers (Fumy *et al.*, 2021). Abandonment results in the same but stronger effects as described above for undergrazing (Löffler *et al.*, 2019; Fumy *et al.*, 2020). Intensive grassland management also leads to homogeneous and dense swards. In such stands, key microhabitats for oviposition (bare ground or taller vegetation that remains during winter) as well as shelter against predators and adverse weather (taller vegetation) are usually lacking (Fumy *et al.*, 2021). Moreover, each mowing event causes direct mortality of grasshopper individuals and, subsequently, facilitates predation through insectivorous vertebrates, since sheltered vegetation has been removed (Wunsch *et al.*, 2012; Buri *et al.*, 2013). Hence, we explain the lower grasshopper densities in general and of threatened species in controls in particular by the high share of abandoned and intensively used plots, respectively. Additionally, we attribute the lack of indicator species of controls in general by the wide gradient in land-use intensity and, hence, strong differences in habitat structure (unpublished data; see also Löffler *et al.*, 2019; Fumy *et al.*, 2020, 2021) although averaged values were often comparable to those of common pastures.

The density of all and of threatened grasshopper species was higher in common pastures compared to control for two vegetation types: grasslands on mineral soil and fens. Both vegetation types occurred on the most productive soils across the studied environmental gradient and were characterized by the highest grazing intensity. By contrast, in controls, only one half and two thirds of the plots, respectively, had a low land-use intensity (mowing once). The remaining plots were either intensively managed (especially in grasslands on mineral soil) or abandoned (particularly in fens). These findings highlight that consistent low-intensity grazing on productive soils favors a high abundance of grasshoppers in general and threatened species in particular. This type of traditional grazing even promotes coexistence of species with very different habitat requirements due to the creation of heterogeneous swards. In grasslands on

mineral soil, the threatened species *C. dorsatus* as well as *P. parallelus*, *R. roeselii*, and *T. subulata* were identified as indicator species of common pastures. *C. dorsatus* is a characteristic of wet grasslands with low land-use intensity; *P. parallelus* occurs in a wide range of different grassland types; *R. roeselii* depends on dense grassland vegetation with an intermediate height, and *T. subulata* colonizes open and wet habitats rich in bare ground (Schlumprecht & Waeber, 2003). By contrast, *C. albomarginatus* was ascertained as the only indicator species of controls, which we attribute to the high share of intensively managed plots in grasslands on mineral soil. Indeed, *C. albomarginatus* is known to belong to the very few grasshopper species that tolerate intensive grassland management (Schlumprecht & Waeber, 2003). In fens, the threatened species *P. montanus* was detected as an indicator species of common pastures; additionally, *C. fuscus* and the threatened species *D. verrucivorus* were indicative for controls. *P. montanus* is a specialized species of wet grasslands and fens that requires a low intensity of land use to establish strong populations (Schlumprecht & Waeber, 2003). By contrast, *C. fuscus* and *D. verrucivorus* depend on taller vegetation, at least in some parts of their habitat (Schlumprecht & Waeber, 2003; Schirmel *et al.*, 2010; Wünsch *et al.*, 2012). Consequently, we explain the occurrence of these two species by the high proportion of abandoned fens in controls.

In contrast to the two more productive vegetation types, in transition mires and raised bogs, the differences in species composition between common pastures and controls were only weak. We attribute this finding to the very slight differences in land use (very low grazing pressure [common pastures] vs. abandonment [control]) and the generally low successional speed within both vegetation types (*cf.* Ellenberg & Leuschner, 2010). Nevertheless, we identified at least one indicator species: the threatened species *M. brachyptera* was indicative for raised bogs in controls. Within raised bogs, the species prefers dense vegetation (Schlumprecht & Waeber, 2003). Indeed, vegetation density near the ground in this vegetation type was higher in controls compared to common pastures. We assume that the abandonment of all control plots and at least weak grazing effects at common pastures were responsible for this difference.

In conclusion, land use had a clear effect on the density of all and of threatened grasshopper species. Traditional low-intensity grazing in common pastures has resulted in open and heterogeneous swards with some bare ground, a low cover of litter and an intermediate vegetation height favoring a high abundance of species in general and threatened species in particular. This is especially true for the two most productive vegetation types, grasslands on mineral soil and fens.

Implications for conservation

Recent studies have shown that traditionally grazed common pastures in southern Bavarian are hotspots of plant and insect diversity, including many rare species (Anthes *et al.*, 2003; Lederbogen *et al.*, 2004; Rosenthal & Lederbogen, 2008; Streitberger *et al.*, 2012; Weking *et al.*, 2013; Helbing *et al.*, 2014; Schwarz & Fartmann, 2021). Our study now highlights that traditional summer cattle grazing in grasslands on mineral soil and fens also enhances grasshopper density in general and density of threatened species in particular.

Grasshoppers are of great functional importance in grassland ecosystems due to their key role as prey for species of higher trophical levels such as insectivorous birds (Samways, 2005; Fartmann *et al.*, 2021). Schwarz *et al.* (2018) showed that the insectivorous Tree Pipit (*Anthus trivialis*) preferred the common pastures in the pre-Alps for breeding. In part, they explained this preference by a higher availability of arthropod food resources in comparison to the surrounding landscape. Our study now provides hard field data that support this assumption.

To promote biodiversity in general and grasshopper densities in particular, we recommend maintaining traditional cattle grazing (stocking capacities: 0.5–2.0 LSU/ha) from May to October in common pastures. Where possible, this grazing regime should also be introduced in the surrounding landscape – however, not at the expense of traditionally managed litter meadows since they are also characterized by highly diverse plant and animal assemblages that are adapted to late mowing (Anthes *et al.*, 2003; Settele *et al.*, 2009; Schwarz & Fartmann, 2021).

Acknowledgments

We are grateful to Wolfgang Kraus (Landratsamt Garmisch-Partenkirchen), Sabine Kraus and Annette Saitner (Landratsamt Bad Tölz-Wolfratshausen), Dieter Frisch (Landratsamt Ostallgäu), and Johannes Wöfl (Landratsamt Weilheim-Schongau) for providing contact data for the members of the grazing associations. Additionally, we would like to thank the grazing associations. We are also grateful to C. Drenker, M. Freienstein, F. Klauer, K. Lebus, E. Notz, and G. Stuhldreher for their support during fieldwork. Lisa Holtmann, Sania Jardine, and Marvin Röthemeyer made valuable comments on an earlier version of the manuscript.

Open access funding enabled and organized by Projekt DEAL.

Disclosure

The authors have no conflicts of interest to declare that are relevant to the content of this article.

Research involving human and animal rights

No grasshoppers were harmed during the course of this project.

References

- Ackermann, W., Balzer, S., Ellwanger, G., Gnittke, I., Kruess, A., May, R., *et al.* (2012) Hotspots der biologischen Vielfalt in Deutschland. Auswahl und Abgrenzung als Grundlage für das Bundesförderprogramm zur Umsetzung der Nationalen Strategie zur biologischen Vielfalt. *Natur und Landschaft*, 87, 289–297.
- Anthes, N., Fartmann, T., Hermann, G. and Kaule, G. (2003) Combining larval habitat quality and metapopulation structure – the key for successful management of pre-alpine *Euphydryas aurinia* colonies. *Journal of Insect Conservation*, 7, 175–185.
- Barnosky, A.D., Matzke, N., Tomiya, S., Wogan, G.O.U., Swartz, B., Quental, T.B. *et al.* (2011) Has the Earth's sixth mass extinction already arrived? *Nature*, 471, 51–57.
- Bartón, M. (2016) Package MuMIn. <https://www.r-project.org> [Accessed 2nd February 2020].
- Bayerisches Geologisches Landesamt (ed.) (1996) Geologische Karte von Bayern 1 : 500 000, 4th edn, München.
- Bazelet, C.S. and Samways, M.J. (2012) Grasshopper and butterfly local congruency in grassland remnants. *Journal of Insect Conservation*, 16, 71–85.
- Bellmann, H. (2006) *Der Kosmos Heuschreckenführer*, Franck-Kosmos Verlags-GmbH & Co. KG, Stuttgart.
- BfN (Bundesamt für Naturschutz) (2012) Landschaftssteckbriefe. <https://www.bfn.de> [Accessed 6th October 2020].
- Brown, K.M. (2006) New challenges for old commons: the role of historical common land in contemporary rural spaces. *Scottish Geographical Journal*, 122, 109–129.
- Buri, P., Arlettaz, R. and Humbert, J.Y. (2013) Delaying mowing and leaving uncut refuges boosts orthopterans in extensively managed meadows: evidence drawn from field-scale experimentation. *Agriculture, Ecosystems & Environment*, 181, 22–30.
- Burnham, K.P. and Anderson, D.R. (2002) *Model Selection and Multimodel Inference. A Practical Information-Theoretic Approach*, Springer, New York.
- Cardoso, P., Barton, P.S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., *et al.* (2020) Scientists' warning to humanity on insect extinctions. *Biological Conservation*, 242, 108426.
- Chytrý, M., Dražil, T., Hájek, M., Kalníková, V., Preislerová, Z., Šibík, J., *et al.* (2015) The most species-rich plant communities in the Czech Republic and Slovakia (with new world records). *Preslia*, 87, 217–278.
- Dengler, J., Janisová, M., Török, P. and Wellstein, C. (2014) Biodiversity of Palaearctic grasslands: a synthesis. *Agriculture, Ecosystems & Environment*, 182: 1–14.
- Dierßen, K. and Dierßen, B. (2008) *Moore: Ökosysteme Mitteleuropas aus geobotanischer Sicht*, Eugen Ulmer, Stuttgart.
- Dirzo, R., Young, H.S., Galetti, M., Ceballos, G., Isaac, N.J.B. and Collen, B. (2014) Defaunation in the Anthropocene. *Science*, 345, 401–406.
- Dufrêne, M. and Legendre, P. (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs*, 67, 345–366.
- DWD (Deutscher Wetterdienst) (2020) Climate Data Center. <https://cdc.dwd.de> [Accessed 6th October 2020].
- EC (European Commission) (2007) The Habitats Directive 92/43/EEC. European Community, Brussels.
- Ellenberg, H. and Leuschner, C. (2010) *Vegetation Mitteleuropas mit den Alpen*, 6th edn, Eugen Ulmer, Stuttgart.
- Fartmann, T., Behrens, M. and Loritz, H. (2008) Orthopteran communities in the conifer-broadleaved woodland zone of the Russian Far East. *European Journal of Entomology*, 105, 673–680.
- Fartmann, T., Jedicke, E., Stuhldreher, G. and Streitberger, M. (2021) *Insektensterben in Mitteleuropa – Ursachen und Gegenmaßnahmen*, Eugen Ulmer, Stuttgart.
- Fartmann, T., Krämer, B., Stelzner, F. and Poniatowski, D. (2012) Orthoptera as ecological indicators for succession in steppe grassland. *Ecological Indicators*, 20, 337–344.
- Feurdean, A., Ruprecht, E., Molnár, Z., Hutchinson, S.M. and Hickler, T. (2018) Biodiversity-rich European grasslands: ancient, forgotten ecosystems. *Biological Conservation*, 228, 224–232.
- Fischer, J., Steinlechner, D., Zehm, A., Poniatowski, D., Fartmann, T., Beckmann, A., *et al.* (2020) *Die Heuschrecken Deutschlands und Nordtirols*. Bestimmen – Beobachten – Schützen, 2nd edn, Quelle & Meyer Verlag, Wiebelsheim.
- Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., *et al.* (2005) Global consequences of land use. *Science*, 309, 570–574.
- Fumy, F., Kämpfer, S. and Fartmann, T. (2021) Land-use intensity determines grassland Orthoptera assemblage composition across a moisture gradient. *Agriculture, Ecosystems & Environment*, 315, 107424.
- Fumy, F., Löffler, F., Samways, M.J. and Fartmann, T. (2020) Response of Orthoptera assemblages to environmental change in a low-mountain range differs among grassland types. *Journal of Environmental Management*, 256, 109919.

- Gardiner, T. and Dover, J. (2008) Is microclimate important for Orthoptera in open landscapes? *Journal of Insect Conservation*, 12, 705–709.
- Gardiner, T., Hill, J. and Chesmore, D. (2005) Review of the methods frequently used to estimate the abundance of Orthoptera in grassland ecosystems. *Journal of Insect Conservation*, 9, 151–173.
- Gepp, J. (2001) Entomologische Relevanz der Fauna-Flora-Habitat-Richtlinie der Europäischen Union. *Entomologica Austriaca*, 1, 7–10.
- Grueber, C.E., Nakagawa, S., Laws, R.J. and Jamieson, I.G. (2011) Multimodel inference in ecology and evolution: challenges and solutions. *Journal of Evolutionary Biology*, 24, 699–711.
- Helbing, F., Blaeser, T.P., Löffler, F. and Fartmann, T. (2014) Response of Orthoptera communities to succession in alluvial pine woodlands. *Journal of Insect Conservation*, 18, 215–224.
- Hothorn, T., Bretz, F. and Westfall, P. (2008) Simultaneous inference in general parametric models. *Biometrical Journal*, 50, 346–363.
- Hrvatin, M. and Perko, D. (2008) Landscape characteristics of common land in Slovenia. *Acta Geographica Slovenica*, 48, 7–32.
- Ingrisch, S. and Köhler, G. (1998) *Die Heuschrecken Mitteleuropas. Die neue Brehm-Bücherei Bd. 629*, Westarp Wissenschaften-Verlagsgesellschaft, Magdeburg.
- IPBES (Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Service) (2019) *Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*, Bonn.
- Kruess, A. and Tschardt, T. (2002) Grazing intensity and the diversity of grasshoppers, butterflies, and trap-nesting bees and wasps. *Conservation Biology*, 16, 1570–1580.
- Lederbogen, D., Rosenthal, G., Scholle, D., Trautner, J., Zimmermann, B. and Kaule, G. (2004) Allmendweiden in Südbayern: Naturschutz durch landwirtschaftliche Nutzung. *Angewandte Landschaftsökologie*, 62, 1–469.
- Löffler, F. and Fartmann, T. (2017) Effects of landscape and habitat quality on Orthoptera assemblages in pre-alpine calcareous grasslands. *Agriculture, Ecosystems & Environment*, 248, 71–81.
- Löffler, F., Poniowski, D. and Fartmann, T. (2016) *Genista* dwarf shrubs as key structures for a Mediterranean grasshopper species on alluvial gravel banks. *Journal of Insect Conservation*, 20, 781–788.
- Löffler, F., Poniowski, D. and Fartmann, T. (2019) Orthoptera community shifts in response to land-use and climate change – lessons from a long-term study across different grassland habitats. *Biological Conservation*, 236, 315–323.
- Marini, L., Fontana, P., Scotton, M. and Klimek, S. (2008) Vascular plant and Orthoptera diversity in relation to grassland management and landscape composition in the European Alps. *Journal of Applied Ecology*, 45, 361–370.
- Nakagawa, S., Johnson, P.C.D. and Schielzeth, H. (2017) The coefficient of determination R^2 and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *Journal of the Royal Society Interface*, 14, 20170213.
- Nakagawa, S. and Schielzeth, H. (2013) A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4, 133–142.
- Oberdorfer, E. (ed.) (1992) *Süddeutsche Pflanzengesellschaften. Fels- und Mauergesellschaften, alpine Fluren, Wasser-, Verlandungs- und Moorgesellschaften*, 3rd edn, Spektrum Akademischer Verlag, Jena, Stuttgart, New York.
- Pliener, T., Hartel, T., Martín-López, B., Beaufoy, G., Bergmeier, E., Kirby, K., et al. (2015) Wood-pastures of Europe: geographic coverage, social-ecological values, conservation management, and policy implications. *Biological Conservation*, 190, 70–79.
- Pliener, T., Höchtl, F. and Spek, T. (2006) Traditional land-use and nature conservation in European rural landscapes. *Environmental Science & Policy*, 9, 317–321.
- Poniowski, D. and Fartmann, T. (2008) The classification of insect communities: lessons from orthoptera assemblages of semi-dry calcareous grasslands in central Germany. *European Journal of Entomology*, 105, 659–671.
- Poniowski, D. and Fartmann, T. (2010) What determines the distribution of a flightless bush-cricket (*Metrioptera brachyptera*) in a fragmented landscape? *Journal of Insect Conservation*, 14, 637–645.
- Poschlod, P. and Braun-Reichert, R. (2017) Small natural features with large ecological roles in ancient agricultural landscapes of Central Europe – history, value, status, and conservation. *Biological Conservation*, 211(B), 60–68.
- Ripple, W.J., Wolf, C., Newsome, T.M., Galetti, M., Alamgir, M., Crist, E., et al. (2017) World scientists' warning to humanity: a second notice. *BioScience*, 67, 1026–1028.
- Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F.S., Lambin, E.F., et al. (2009) A safe operating space for humanity. *Nature*, 461, 472–475.
- Rosenthal, G. and Lederbogen, D. (2008) Response of the clonal plant *Apium repens* (Jacq.) Lag. to extensive grazing. *Flora*, 203, 141–151.
- Samways, M.J. (2005) *Insect Diversity Conservation*, Cambridge University Press, Cambridge.
- Sánchez-Bayo, F. and Wyckhuys, K.A.G. (2019) Worldwide decline of the entomofauna: a review of its drivers. *Biological Conservation*, 232, 8–27.

- Scherer, G., Löffler, F. and Fartmann, T. (2021) Abandonment of traditional land use and climate change threaten the survival of an endangered relict butterfly species. *Insect Conservation and Diversity*, 14, 556–567.
- Schirmel, J., Blindow, I. and Fartmann, T. (2010) The importance of habitat mosaics for Orthoptera (Caelifera and Ensifera) in dry heathlands. *European Journal of Entomology*, 107, 129–132.
- Schlumprecht, H. and Waeber, G. (2003) *Heuschrecken in Bayern*, Eugen Ulmer, Stuttgart.
- Schwarz, C. and Fartmann, T. (2021) Conservation of a strongly declining butterfly species depends on traditionally managed grasslands. *Journal of Insect Conservation*, 25, 255–271.
- Schwarz, C., Trautner, J. and Fartmann, T. (2018) Common pastures are important refuges for a declining passerine bird in a pre-alpine agricultural landscape. *Journal of Ornithology*, 159, 945–954.
- Sergeev, M.G. (2021) Distribution patterns of grasshoppers and their kin over the Eurasian steppes. *Insects*, 12, 77.
- Settele, J., Steiner, R., Reinhardt, R., Feldmann, R. and Hermann, G. (2009) *Schmetterlinge: die Tagfalter Deutschlands*, Eugen Ulmer, Stuttgart.
- Sheather S.J. (2009) *A Modern Approach to Regression with R*. Springer, New York.
- Sliacka, A., Krištin, A. and Nad'ó, L. (2013) Response of Orthoptera to clear-cuts in beech forests. *European Journal of Entomology*, 110, 319–326.
- Stork, N.E. (2018) How many species of insects and other terrestrial arthropods are there on Earth? *Annual Review of Entomology*, 63, 31–45.
- Stoutjesdijk, P. and Barkman, J.J. (1992) *Microclimate, vegetation and fauna*, Opulus Press, Upsala.
- Streitberger, M., Hermann, G., Kraus, W. and Fartmann T. (2012) Modern forest management and the decline of the Woodland Brown (*Lopinga achine*) in Central Europe. *Forest Ecology and Management*, 269, 239–248.
- Succow, M. and Jeschke, L. (1990) *Moore in der Landschaft*, Urania-Verlag, Leipzig.
- Thomas, C.D., Franco, A.M.A. and Hill, J.K. (2006) Range retractions and extinction in the face of climate warming. *Ecology and Evolution*, 21, 415–416.
- Thomas, J.A., Telfer, M.G., Roy, D.B., Preston, C.D., Greenwood, J.J.D., Asher, J., et al. (2004) Comparative losses of British butterflies, birds, and plants and the global extinction crisis. *Science*, 303, 1879–1881.
- Thumm, U., Ehrmann, O. and Wagner, F. (2005) Spatial variation of grazing effects in cooperatively used pastures in the pre-Alpine region of Southern Germany. *Integrating Efficient Grassland Farming and Biodiversity*, 10, 356–359.
- Uchida, K. and Ushimaru, A. (2014) Biodiversity declines due to abandonment and intensification of agricultural lands: patterns and mechanisms. *Ecological Monographs*, 84, 637–658.
- Voith, J., Beckmann, A., Sachteleben, J., Schlumprecht, H. and Waeber, G. (2016) Rote Liste und Gesamtartenliste der Heuschrecken (Saltatoria) Bayerns. *Bayerisches Landesamt für Umwelt (LfU)*. https://www.lfu.bayern.de/natur/rote_liste_tiere/2016/index.htm [Accessed 1st April 2020].
- Wagner, D.L. (2020) Insect declines in the Anthropocene. *Annual Review of Entomology*, 65, 457–480.
- Wallis de Vries, M. and Poschlod, P. (2002) Challenges for the conservation of calcareous grasslands in Northwestern Europe. *Biological Conservation*, 104, 265–273.
- Wünsch, Y., Schirmel, J. and Fartmann, T. (2012) Conservation management of coastal dunes for Orthoptera has to consider oviposition and nymphal preferences. *Journal of Insect Conservation*, 16, 501–510.
- Wettstein, W. and Schmid, B. (1999) Conservation of arthropod diversity in montane wetlands: effect of altitude, habitat quality and habitat fragmentation on butterflies and grasshoppers. *Journal of Applied Ecology*, 36, 363–373.
- Weking, S., Hermann, G. and Fartmann, T. (2013) Effects of mire type, land use and climate on a strongly declining wetland butterfly. *Journal of Insect Conservation*, 17, 1081–1091.

Manuscript received October 22, 2021

Final version received March 18, 2022

Accepted March 22, 2022

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1 Results of Spearman rank correlations (rs) of environmental variables.

Table S1 Results of Spearman rank correlations (r_s) of environmental variables. Only variables with strong inter-correlations ($|r_s| \geq 0.5$) are displayed and highlighted in bold type. ‘No. of cow droppings’ and ‘Vegetation density (VD) in 0–5 cm height’ were excluded from GLMM (cf. Table 4). For further information, cf. Section Statistical analysis. $**P \leq 0.01$, $***P \leq 0.001$.

Variable	Cover of Poaceae	VD 0–5 cm	No. cow droppings
Cover of shrubs	–0.44**	–0.52***	–0.54***
Cover of Poaceae	/	0.52***	0.63***
VD 0–5 cm	.	/	0.40**